

Permanent confinement in compact QED₃ with fermionic matter

Igor F. Herbut and Babak H. Seradjeh

Department of Physics, Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6

We argue that the compact three dimensional electrodynamics with massless relativistic fermions is always in the confined phase, in spite of the bare interaction between the magnetic monopoles being rendered logarithmic by fermions. The effect is caused by screening by other dipoles, which transforms the logarithmic back into the Coulomb interaction at large distances. Possible implications for the chiral symmetry breaking for fermions are discussed, and the global phase diagram of the theory is proposed.

It is well known that the pure compact electrodynamics in three dimensions (3D) is always in the confined phase, i. e. in the phase where magnetic monopoles form a neutral plasma of free magnetic charges [1]. When the gauge field is coupled to matter, however, new possibilities emerge. In particular, Abelian gauge fields coupled to massless relativistic fermions (cQED₃) arise naturally in several theories of high temperature superconductivity [2], [3]. cQED₃ also represents maybe the simplest theory that should contain the physics of confinement and chiral symmetry breaking, and as such it may be used as a valuable toy model for the QCD. It has been argued that coupling to massless relativistic fermions turns the interaction between monopoles from $1/x$ to $-\ln(x)$ at large distances x [4], [5], [6], [7], so that the deconfined phase with bound monopole-antimonopole pairs may become stable at low (effective) temperatures, in close analogy to the Kosterlitz-Thouless (KT) transition. If so, this would suggest that, at least for some values of coupling constants, compactness of the gauge field may be neglected at large distances, and that the results pertaining to the continuum QED₃ [8] are likely to remain valid. In particular, the chiral symmetry breaking instability of the continuum QED₃ that has been linked to antiferromagnetism [9] in cuprates might then be expected to survive essentially intact.

In this paper we consider the effects of finite density of monopole-antimonopole pairs (dipoles) on the putative, KT-like, confinement-deconfinement transition in the cQED₃. We find that, in stark contrast to two dimensions (2D), screening by logarithmically interacting magnetic charges bound to dipoles in 3D always alters the *form* of the interaction between the charges, from $-\ln(x)$ at large distance, to $1/x$. We demonstrate this first by an elementary electrostatic argument, and then support it by a controlled momentum shell renormalization group on the equivalent sine-Gordon theory. As a result, largest dipoles are always unbound, and would be expected to provide linear confinement of electric charges. Finally, we argue that the absence of the phase transition in the cQED₃ in terms of magnetic monopoles should imply the same in terms of fermions. Relying on the existing numerical results and qualitative arguments, we suggest that the phase of free monopoles corresponds to the (confined) phase of broken chiral symmetry for fermions, and that the (deconfined) chirally symmetric phase exists

only in the continuum limit, and above a certain number of fermion species [8]. The conjectured phase diagram is presented in Fig. 1.

We are interested in the cQED₃ defined on a cubic lattice:

$$S = S_F(\chi, \theta_\mu) - \beta \sum_{x, \mu, \nu} \cos(F_{\mu\nu}), \quad (1)$$

where $F_{\mu\nu} = \epsilon_{\mu\nu\beta} \epsilon_{\beta\rho\sigma} \Delta_\rho \theta_\sigma$ is the lattice version of the electromagnetic tensor, and the coupling to fermions on a lattice may be written as

$$S_F = \frac{1}{2} \sum_{x, \nu} \sum_{i=1}^{N/2} \eta_\nu(x) [\bar{\chi}_i(x) e^{i\theta_\nu(x)} \chi_i(x + \hat{\nu}) - \bar{\chi}_i(x + \hat{\nu}) e^{-i\theta_\nu(x)} \chi_i(x)], \quad (2)$$

using the standard staggered fermions. Here, x denotes the lattice sites and $\hat{\nu}$ the direction of the links. In the continuum, S_F corresponds to N species of four component Dirac fermions [10].

The cQED₃ has been previously studied numerically in [11]. Here, to make progress analytically and following [7], we approximate (1) with

$$S \approx \frac{1}{2} \sum_{x, \nu, \mu} [F_{\mu\nu}(x) - 2\pi n_{\mu\nu}(x)] \left(\beta + \frac{N}{16|\Delta|} \right) [F_{\mu\nu}(x) - 2\pi n_{\mu\nu}(x)], \quad (3)$$

where $1/|\Delta|$ should be understood as an inverse of the square root of the lattice gradient squared. The term proportional to the coupling β in Eq. (3) is the standard Villain approximation to the second term in (1). The form of the term proportional to N in (3) is motivated by the known continuum limit of the fermion polarization: neglecting compactness of a_μ by setting $n_{\mu\nu}(x) \equiv 0$ would give the standard one-loop Gaussian action in the continuum, with the correct coefficient [8]. Just like in the $\sim \beta$ term, here we also retained only the leading power of fields and their lattice derivatives. Eq. (3) may be understood as the quadratic approximation to cQED₃ in Eq. (1). It leads to the expected logarithmic interaction between the magnetic monopoles, and therefore appears to contain the essential physics of cQED₃ we wish to address.




FIG. 1: The proposed phase diagram of the cQED₃. True phase transition between the chirally symmetric (CS) and chiral symmetry broken (CSB) phases of fermions occurs only in the continuum limit ($\beta \rightarrow \infty$). Monopoles are in the plasma phase at all $\beta < \infty$ and $N < \infty$. Dashed line marks a crossover that corresponds to the $N_{\text{cross}}(\beta)$, as discussed in the text.

Standard duality transformations [12] on (3) lead to the sine-Gordon action in the continuum

$$S_{\text{SG}} = \int d^3\vec{x} \left[\frac{(\vec{\nabla}\Phi(\vec{x}))|\vec{\nabla}|(\vec{\nabla}\Phi(\vec{x}))}{\pi^2(N + 16\beta|\vec{\nabla}|)} - 2y \cos \Phi(\vec{x}) \right], \quad (4)$$

where $y \ll 1$ is the fugacity of monopoles, and $\vec{x}$ is now the position vector in the continuum. In presence of fermions ($N \neq 0$), at small momenta one can expand in the original gauge coupling β in Eq. (4), which becomes a coefficient of the term quartic in derivatives, and thus irrelevant by power counting. We may therefore safely set it to zero, and return to its (non-universal) effects on the phase diagram later. We also find it useful to have an alternative representation of the partition function defined by S_{SG} ,

$$Z = \sum_{n=0}^{\infty} \frac{y^n}{n!} \int \prod_{i=1}^n d^3\vec{x}_i \exp \left[-\frac{1}{2T} \sum_{i,j} q_i q_j V(|\vec{x}_i - \vec{x}_j|) \right], \quad (5)$$

in terms of unit magnetic charges $q_i = \pm 1$ at an effective temperature $T = 4/N + O(\beta)$, interacting at large distance via $V(x) = -\ln(x\Lambda)$. Λ is the ultraviolet cutoff implicit in Eq. (4). The sine-Gordon field in Eq. (4) is then $\langle \Phi(\vec{x}) \rangle = (i/T) \sum_i q_i \langle V(|\vec{x} - \vec{x}_i|) \rangle$, where $\langle \dots \rangle$ denotes averaging with respect to Z .

The excess of free energy due to a *single* isolated

monopole in the sample of a linear size L is therefore

$$\Delta F = \frac{1}{2} \ln L - T \ln L^3, \quad (6)$$

so that for $T < 1/6$ monopoles would be expected to be bound in pairs. This corresponds to a “critical” number of fermion components $N = 24$ [4], [6], [7]. This argument neglects the effect of other dipoles, however. In the standard KT transition their presence does not change the critical temperature obtained by the simple energy-entropy argument, since the KT fixed point lies at zero fugacity [13]. We show next that the situation in 3D is radically different.

Let us assume a distribution of charges $\rho(\vec{x})$ interacting via $-\ln(x)$ interactions in 3D, located in a region of a finite size R . At a distance $x \gg R$ we can write the interaction as

$$V(\vec{x}) = \int d^3\vec{x}' \rho(\vec{x}') \left(-\ln|\vec{x}| + \frac{\vec{x} \cdot \vec{x}'}{x^2} + \dots \right) \quad (7)$$

in spirit of multipole expansion [14]. Thus, for a medium with a monopole density $\rho(x)$ and a dipole moment density $\vec{P}(\vec{x}) = \rho(\vec{x})\vec{x}$, the potential is given by

$$V(\vec{x}) = \int d^3\vec{x}' (-\ln|\vec{x} - \vec{x}'|) [\rho(\vec{x}') - \vec{\nabla}' \cdot \vec{P}(\vec{x}') + \dots]. \quad (8)$$

The energy of a small dipole with a moment $\vec{P} = q\vec{r}$ ($q = 1$) at $\vec{x}$ in a weak potential $V(\vec{x})$ is still $\vec{P} \cdot \nabla V(\vec{x})$. Assuming non-interacting dilute dipoles at a temperature T then gives the average dipole moment at $\vec{x}$

$$\langle \vec{P}(\vec{x}) \rangle = \frac{\langle r^2 \rangle}{3T} (-\nabla V(\vec{x})). \quad (9)$$

Since the density of dipoles is proportional to y^2 , the polarizability of the medium is thus

$$\chi \sim \frac{\langle r^2 \rangle}{3T} y^2, \quad (10)$$

where the constant of proportionality depends on the precise form of the short distance regularization of the logarithmic interaction.

Writing Eq. (8) in Fourier space and combining Eqs. (9) and (10) into $\vec{P}(\vec{q}) = -i\chi\vec{q}V(\vec{q})$, we finally obtain the interaction due to an external point charge Q in presence of a finite density of dipoles

$$V(\vec{q}) = \frac{Q}{|\vec{q}|^3 + \chi q^2}, \quad (11)$$

or, in real space

$$V(\vec{x}) = \frac{1}{4\pi\chi} \frac{1}{x} + O\left(\frac{1}{x^2}\right). \quad (12)$$

This implies that in presence of other dipoles the first (energy) term in Eq. (6) scales like $1/L$, so that the entropy always dominates, and the largest dipoles should




FIG. 2: The flow diagram in the $\hat{y}^2$ - $\hat{a}$ plane for $N < N_{\text{cross}}$ (a), and $N > N_{\text{cross}}$ (b). The non-trivial fixed point in (a) moved to lower right quadrant (not shown) in (b).

dissociate into free monopoles. The ‘dielectric’ phase of logarithmically interacting monopoles in 3D thus, unlike in 2D, appears destabilized by dipole screening, as anticipated in [15]. A similar conclusion has apparently also been drawn before in the study of instantons in quantum antiferromagnets [16].

The renormalization group treatment of the equivalent sine-Gordon theory closely parallels the electrostatic consideration. Integrating out the modes of $\Phi(\vec{x})$ in Eq. (4) with momenta $\Lambda/b < q < \Lambda$ one finds that the original couplings are renormalized as

$$N(b) = N + O(y^3), \quad (13)$$

$$y(b) = b^3 y e^{-\frac{1}{2}G_{>}(0)} + O(y^3), \quad (14)$$

$$a(b) = b \left[a + \frac{1}{3} y^2 e^{-G_{>}(0)} \int d^3x x^2 (e^{G_{>}(x)} - 1) \right] + O(y^3), \quad (15)$$

where a is the coefficient in front of the $(\vec{\nabla}\Phi(\vec{x}))^2$ term in S_{SG} , Eq. (4). Such a term is initially absent, but gets *generated* to the second order in y . Formally this is just what one expects, since there is no particular symmetry in the Eq. (4) that would prohibit its appearance. Physically, as we have seen, this simply expresses the screening effect of smaller dipoles onto the interaction between the monopoles in a larger dipole. This term was effectively overlooked in the previous work on cQED₃ [4], [5], [6], [7], where the authors consequently arrived at a different final result.

The correlation function appearing in the recursion re-

lations (14) and (15) is

$$G_{>}(\vec{x}) = \int_{\Lambda/b < |\vec{q}| < \Lambda} \frac{d^3\vec{q}}{(2\pi)^3} \frac{\pi^2 N e^{i\vec{q}\cdot\vec{x}}}{2|\vec{q}|^3 + \pi^2 N a q^2}, \quad (16)$$

and requires some care in evaluation. Assuming a low cutoff, or the effective temperatures $T\Lambda^3 \ll y$, we may rewrite it as

$$G_{>}(\vec{x}) = \frac{\pi^2 N}{2\Lambda + \pi^2 N a} \int \frac{d^3\vec{q}}{(2\pi)^3} \frac{e^{i\vec{q}\cdot\vec{x}}}{q^2} \left(\frac{q^2}{q^2 + (\Lambda/b)^2} - \frac{q^2}{q^2 + \Lambda^2} \right), \quad (17)$$

where the integration is now unconstrained, but we have introduced a smoothing function $q^2/(q^2 + \Lambda^2)$ instead of the sharp cutoff [17]. We have checked that when applied to the standard sine-Gordon problem in 2D, our regularization scheme gives the correct anomalous dimension at the KT transition. With this regularization we find

$$G_{>}(\vec{x}) = \frac{1}{4\pi} \frac{\pi^2 N \ln(b)}{2 + \pi^2 N \hat{a}} e^{-x\Lambda} + O((\ln b)^2), \quad (18)$$

and therefore

$$\dot{N} = 0 + O(\hat{y}^3), \quad (19)$$

$$\dot{\hat{y}} = \left[3 - \frac{\pi N}{8(2 + \pi^2 N \hat{a})} \right] \hat{y} + O(\hat{y}^3), \quad (20)$$

$$\dot{\hat{a}} = \hat{a} + \frac{8\pi^2 N}{2 + \pi^2 N \hat{a}} \hat{y}^2 + O(\hat{y}^3), \quad (21)$$

where $\hat{y} = y/\Lambda^3$ and $\hat{a} = a/\Lambda$, are the dimensionless couplings, and $\dot{z} \equiv dz/d \ln b|_{b=1}$.

Two important features of the flow equations (19)-(21) should be noted: 1) the relevant coupling a , even if absent initially, becomes generated at lower cutoffs, and 2) N is marginal to $O(y^3)$. We suspect N to be an exactly marginal coupling, since the action for slow modes should be analytic in $(q/\Lambda)^2$, so the coefficient of the non-analytic $|q|^3$ term can not get renormalized to any order [18].

The flow in the $\hat{y}^2$ - $\hat{a}$ plane for $N < N_{\text{cross}}$ and $N > N_{\text{cross}}$, $N_{\text{cross}} = 48/\pi$, is depicted in Fig. 2. Starting from $\hat{a} = 0, \hat{y} = \hat{y}_0$, the fugacity in the former case monotonically increases as $b \rightarrow \infty$, while in the latter it begins to increase only for $b > b^*$, with $b^*(\hat{y}_0) \approx 1/(96\pi\hat{y}_0)$ at $N \gg 1$. Nevertheless, since $\hat{y} = 0$ is an invariant line under the scale transformations, fugacity must increase at large enough b for any initial value. We interpret this upward flow of the fugacity as an indication that monopoles are always in the plasma phase.

Interestingly, massless fermions still make themselves felt in such a plasma, since the non-analytic $|\vec{\nabla}|^3$ term in Eq. (4) translates into a power-law behavior of the screened interaction at long distances. Instead of the expected Debye-Hückel exponential decay, in the quadratic approximation to (4) we find $V_{\text{scr}}(x) \approx -6/(\pi^4 y^2 N |\vec{x}|^6)$ at large x . The reader may recall a similar phenomenon

of Friedel oscillations in metals, where a non-analyticity due to the Fermi surface produces a power-law behavior of the screened interaction as well. In the present case the sign of the above power-law term indicates *overscreening*, presumably due to the extremely long range of the bare (logarithmic) interaction between monopoles.

Let us next turn to possible implications of our findings for fermions. First, the dimensionless coupling $\beta = 1/(e^2 l)$, where e is the dimensionfull charge, and l the lattice spacing. The lattice gauge field $\theta_\mu(x) = elA_\mu(x)$, where $A_\mu(x)$ is the gauge field in the continuum. The continuum limit is achieved by keeping e constant and taking $l \rightarrow 0$. In other words, $\beta \rightarrow \infty$ should naively correspond to the continuum QED₃. In this theory chiral symmetry for fermions is expected to be broken, and the condensate $\langle \bar{\Psi}\Psi \rangle \neq 0$, for $N < N_{\text{ch}}$, where N_{ch} is of order unity [8], [19], [20]. On the other hand, for $\beta < \infty$ we argued that once fermions are integrated out, the theory has no phase transition. If correct, this implies that, had we proceeded in reverse and integrated out everything but the fermions, we would have had to find them in a single phase at any finite β . Relying on numerical evidence that free monopoles seem to *enhance* chiral symmetry breaking [11], we propose that it is the phase with broken chiral symmetry that survives a finite β .

It was found previously [21] that the critical N for chi-

ral symmetry breaking at $\beta = 0$ is significantly larger than N_{ch} in the continuum limit. We expect that for $\beta = 0$, there is no true phase transition in the cQED₃, just the crossover occurring at N_{cross} . Making the simplest assumption that β always stays irrelevant, the same would hold at any $\beta < \infty$, but with the crossover line shifted to $N_{\text{cross}}(\beta) = N_{\text{cross}} - 16\beta\Lambda + O((\beta\Lambda)^2)$. The conjectured phase diagram that summarizes this discussion is presented in Fig. 1. $\langle \bar{\Psi}\Psi \rangle \neq 0$ everywhere, except on the line $1/\beta = 0$, $N > N_{\text{ch}}$.

To conclude, we argued that magnetic monopoles are always free in the cQED₃ with gapless relativistic fermions, and the theory is thus expected to be permanently confining. We proposed the phase diagram, and conjectured that chiral symmetry for fermions is always spontaneously broken. Our results seem to be in qualitative agreement with the early numerical results of ref. [22]. They can also be generalized to the case with non-relativistic fermions with Fermi surface [23]. Although we have studied only the U(1) theory coupled to fermions in this paper, similar arguments should be applicable to the case of critical scalar field [7], [24].

We thank F. Nogueira and A. Sudbo for useful discussions, and S. Sachdev for remarks that in part stimulated this work. This work was supported by NSERC of Canada and the Research Corporation.

-
- [1] A. M. Polyakov, Nucl. Phys. **B120**, 429 (1977).
 - [2] See D. H. Kim and P. A. Lee, Ann. Phys. (N. Y.) **272**, 130 (1999) for a discussion of the compact gauge-theory of the $t - J$ model.
 - [3] See I. F. Herbut, Phys. Rev. B **66**, 094504 (2002) for the gauge theory of the fluctuating d-wave superconductors.
 - [4] L. B. Ioffe and A. I. Larkin, Phys. Rev. B **39**, 8988 (1989).
 - [5] J. B. Marston, Phys. Rev. Lett. **64**, 1166 (1990).
 - [6] X.-G. Wen, Phys. Rev. B **65**, 165113 (2002).
 - [7] H. Kleinert, F. S. Nogueira, A. Sudbo Phys. Rev. Lett. **88**, 232001 (2002); preprint hep-th/0209132.
 - [8] See, P. Maris, Phys. Rev. D **54**, 4049 (1995) for a review.
 - [9] I. F. Herbut, Phys. Rev. Lett. **88**, 047006 (2002).
 - [10] C. Burden and A. N. Burkitt, Europhys. Lett. **3**, 545 (1987).
 - [11] H. R. Fiebig and R. M. Woloshyn, Phys. Rev. D **42**, 3520 (1990).
 - [12] M. B. Einhorn and R. Savit, Phys. Rev. D **17**, 2583 (1979).
 - [13] J. M. Kosterlitz, J. Phys. C **7**, 1046 (1974).
 - [14] J. D. Jackson, *Classical Electrodynamics* (John Wiley and Sons, Inc., 1975), Ch. 4.
 - [15] S. Sachdev and K. Park, Ann. Phys. (N. Y.) **298**, 58 (2002).
 - [16] G. Murthy, Phys. Rev. Lett. **67**, 911 (1991).
 - [17] T. Ohta and D. Jasnow, Phys. Rev. B **20**, 139 (1979).
 - [18] I. F. Herbut, Phys. Rev. Lett. **87**, 137004 (2001).
 - [19] T. Appelquist, A. G. Cohen, and M. Schmaltz, Phys. Rev. D **60**, 045003 (1999).
 - [20] S. J. Hands, J. B. Kogut, and C. G. Strouthos, Nucl. Phys. **B645**, 321 (2002).
 - [21] E. Dagotto, J. B. Kogut, and A. Kocić, Nucl. Phys. **B334**, 279 (1990).
 - [22] V. Azcoiti and X.-Q. Luo, Mod. Phys. Lett. A **8**, 3635 (1993).
 - [23] I. F. Herbut, B. H. Seradjeh, S. Sachdev, and G. Murthy, preprint cond-mat/0306537.
 - [24] M. N. Chernodub, I.-M. Ilgenfritz, and A. Schiller, Phys. Lett. **B547**, 269 (2002).